

## Effect of generalized thermoelasticity materials with memory

Arup Baksi<sup>†</sup>

*Umes Chandra College 13, Surya Sen Street, Calcutta-700012, India*

Bidyut Kumar Roy<sup>‡</sup>

*Department of Mathematics, Vivekananda College, 269, D. H. Road, Calcutta-700063, India*

Rasajit Kumar Bera<sup>‡‡</sup>

*Heritage Institute of Technology, Chowbaga Road, Anandapur, P.O. East Kolkata Township, Kolkata-700107, India*

*(Received November 10, 2004, Accepted December 20, 2006)*

**Abstract.** Many works have been done in classical theory of thermoelasticity in materials with memory by researchers like Nunziato, Chen and Gurtine and many others. No work is located in generalized thermoelasticity regarding materials with memory till date. The present paper deals with the wave propagation in materials with memory in generalized thermoelasticity. Plane progressive waves and Rayleigh waves have been discussed in details. In the classical theory of heat conduction it was observed that heat propagates with infinite speed. This paradox has been removed in the present discussion. The set of governing equations has been developed in the present analysis. The results of wave velocity and attenuation coefficient corresponding to low and high frequency have been obtained. For thermal wave the results show appreciable differences with those in the usual thermoelasticity theory.

**Keywords:** generalized thermoelasticity; plane progressive waves; Rayleigh waves.

---

### 1. Introduction

In the mechanics of continuous media, a material is said to have hereditary characteristics or memory if its behavior at time  $t$  is specified by the past experience of the body upto time  $t$ . The theory of heat conduction in materials with memory has drawn the attention of many researchers. The motivation was provided by an unpleasant feature of the classical linear theory of heat conduction viz., that a thermal disturbance produced at some point in the body felt instantaneously at all other points, this contradicts the relativistic principle that energy cannot be propagated at

---

<sup>†</sup> E-mail: [arup\\_baksi2003@yahoo.co.in](mailto:arup_baksi2003@yahoo.co.in)

<sup>‡</sup> E-mail: [mht\\_biro@yahoo.co.uk](mailto:mht_biro@yahoo.co.uk)

<sup>‡‡</sup> Professor, Ph.D., Corresponding author, E-mail: [rasajit@yahoo.com](mailto:rasajit@yahoo.com)

speeds exceeding the velocity of light. Gurtin and Pipkin (1968) first established a general non-linear theory of heat conduction in rigid materials with memory for which thermal disturbances propagate with finite speed. They assumed that the response functional viz., the entropy, free energy and heat-flux depend on the present value of the temperature and the integrated histories of the temperature and the temperature gradient.

Nunziato (1971) considered a slightly different memory theory of heat-conduction. He assumed the response functional to depend on the histories upto the present time of the temperature and the temperature gradient. In his theory heat conduction depends also on the present value of the temperature gradient so that Fourier's law of heat conduction is obtained as a particular case, if  $k(0)$ , the instantaneous conductivity, is non-zero. On the other hand, if  $k(0) = 0$ , Nunziato's heat conduction equation agrees with that of Gurtin and Pipkin. Chen and Gurtin (1970) extended the theory presented by Gurtin and Pipkin to deformable media. They started with the constitutive assumptions that the response functional viz., the stress, entropy, free energy and heat-flux depend on the present values of the temperature and the deformation gradient and the integrated histories of the deformation gradient, temperature and the temperature gradient. On the other hand, McCarthy (1970) developed a theory of thermo-mechanical materials with memory on the assumption that the response functionals are dependent on the present values of the temperature and the deformation gradient, histories of the deformation gradient and the temperature and the integrated history of the temperature gradient. Chakraborty (1976), however, assumed the response functionals to be dependent on the present values of the temperature and the deformation gradient and the histories of the temperature, deformation gradient and the temperature gradient. His assumptions are thus less restrictive than those of Chen and Gurtin and of McCarthy. He extended Nunziato's theory of heat conduction to deformable bodies and deduced the generalized stress-strain relations and the coupled equation of heat conduction for materials with memory and obtained the linearised forms of these equations. The stress-strain relations are similar to those of a linear thermo-visco-elastic solid given by Eringen (1967, p.367-368). The heat conduction equation for a thermo-visco-elastic solid is the same as that for the thermo-elastic solid Eringen (1967). It is, thus, observed that the stress-strain relations and the heat conduction equation obtained by Chakraborty (1976) are more general than those of others in the sense that, the equations of classical thermoelasticity as well as those of classical thermo-visco-elasticity can all be obtained as particular cases. The stress-strain relations and the coupled equation of heat conduction in the linear coupled thermo-elasticity theory given by Chadwick (1960) for thermoelastic materials (Müller 1967) can be obtained as a particular case from those deduced by Chakraborty. This is to be expected, since for "thermoelastic" materials the response functionals depend only on the present values of the temperature, temperature gradient, and the deformation gradient. Hence "thermoelastic" materials are special cases of "thermoelastic materials with memory".

Chen, Amos, Nunziato and McCarthy have studied propagation of thermoelastic waves. Chen (1969) studied the amplitude variation of the temperature rate waves of arbitrary form, assuming that the constitutive relations of the material are given by the linearised theory derived by Gurtin and Pipkin (1968). Amos and Chen (1970) obtained the speed of propagation of one dimensional thermal waves, while Nunziato (1971) determined the attenuation and the speed of plane temperature rate waves. McCarthy (1970) studied the propagation of "first order" waves using his own derived equations (McCarthy 1970).

The recent works in this line by Bhattacharyya (1991), Matinez and Quintanilla (1995), Chirita and Quintanilla (1997), Hongjun (2001), Iesan and Quintanilla (2002), Maria Grazia and Yuming.

Qin (2002), Maria Grazia and Federico (2003), Maria Grazia (2003), Zhou *et al.* (2003), Maria Grazia and Elena (2004) may be of worth mentioning.

In the present paper we study plane progressive thermoelastic waves and Rayleigh waves in generalized thermoelasticity in a general form using the more general equations deduced by Chakraborty (1976). As in the usual coupled thermoelasticity theory, it is observed in the present analysis that there exist two types of propagating plane waves, one is the quasi-elastic wave and the other is the quasi-thermal wave (Chadwick 1960, p.283). Both the waves, however, exhibit dispersion and attenuation. The modified elastic wave speed approaches the “instantaneous” value of the classical compressional wave velocity at high frequencies and its “equilibrium” value modified by the coupling factor at low frequencies. Its attenuation coefficient tends to a constant value at large frequencies and is proportional to the square of the frequency at low frequencies. Similar features are observed in the usual coupled thermoelasticity theory. But a new feature, exhibited by purely elastic waves in materials with memory, is that these waves are subject to both dispersion and attenuation. The purely elastic wave speed approaches the “instantaneous” value of the classical compressional wave velocity at high frequencies and its equilibrium value at low frequencies. The attenuation coefficient is proportional to the square of the frequency at low frequencies and tends to a constant value at high frequencies.

It is also observed that for thermal waves both the velocity and the attenuation coefficient are proportional to the square root of the frequency at low and high frequencies.

Finally, the dispersion equation for Rayleigh waves has also been obtained in the present study. It has a form analogous to that in the usual linear coupled thermoelasticity theory.

## 2. Formulation of equation

Following the analysis of Dhaliwal and Sing (1980), Lord and Shulman (1967) and Green and Lindsay (1972) in generalized thermoelasticity, the stress-strain-temperature relations and heat conduction equations are written below:

Assuming isotropy and linearity the equation of heat conduction or energy equation can be written as

$$(i) \quad \alpha(0)(\dot{\theta} + t_0 \ddot{\theta}) + \int_0^\infty \alpha'(s)[\dot{\theta}(\vec{x}, t-s) + t_0 \ddot{\theta}(\vec{x}, t-s)]ds + \eta(0)div[\vec{u}(\vec{x}, t) + \delta_{ik}t_0 \ddot{u}(\vec{x}, t)] \\ + \int_0^\infty \eta'(s)div[\vec{u}(\vec{x}, t-s) + \delta_{ik}t_0 \ddot{u}(\vec{x}, t-s)]ds = k(0)\nabla^2\theta(\vec{x}, t) + \int_0^\infty k'(s)\nabla^2\theta(\vec{x}, t-s)ds \quad (1)$$

(ii) Stress-Strain-temperature relations are

$$\sigma_{ij}(\vec{x}, t) = [\lambda(0)e(\vec{x}, t) - \beta(0)\{\theta(\vec{x}, t) + t_1\dot{\theta}(\vec{x}, t)\}\delta_{ij} + 2\mu(0)e_{ij}(\vec{x}, t)] \\ + \int_0^\infty [\lambda'(s)e(\vec{x}, t-s) - \beta'(s)\{\theta(\vec{x}, t-s) + t_1\dot{\theta}(\vec{x}, t-s)\}\delta_{ij} + 2\mu'(s)e_{ij}(\vec{x}, t-s)]ds \quad (2)$$

(iii) Equations of motion are

$$\sigma_{ij,j}(\vec{x}, t) = \rho \ddot{\vec{u}}(\vec{x}, t) \quad (3)$$

(iv) Strain-displacement relations are

$$e_{ij} = \frac{1}{2}(u_{i,j} + u_{j,i}) \quad i, j = 1, 2, 3 \quad (4a)$$

In Eqs. (1) and (2),  $\alpha(s)$  and  $k(s)$  are respectively the energy-temperature relaxation function and the heat-conduction relaxation function, which in the case of “thermoelastic” materials correspond to specific heat and conductivity respectively.  $\lambda(s)$  and  $\mu(s)$  may be called the Lamé relaxation functions corresponding to the usual Lamé constants.  $\beta(s)$  and  $\eta(s)$  are the relaxation functions corresponding to  $\beta$  and  $\eta$  where  $\beta$  is the coefficient of the temperature deviation in the stress-strain relation, and  $\eta$  is the coupling constant.  $\theta$  denotes temperature deviation and  $\vec{u}(\vec{x}, t)$  is the displacement vector.

It should be noted that the relations

$$\begin{aligned} \beta(0) &= [3\lambda(0) + 2\mu(0)]\alpha_T \\ \eta(0) &= T_0\beta(0) \end{aligned} \quad (4b)$$

where  $\alpha_T$  is the coefficient of linear expansion and  $T_0$  is the reference temperature, hold though they are not valid for other values of the arguments (they are valid for “thermoelastic” materials when  $\alpha(s)$ ,  $\beta(s)$ ,  $\lambda(s)$ ,  $k(s)$ ,  $\eta(s)$ ,  $\mu(s)$  are all constants). If the relaxation functions  $\alpha(s)$ ,  $\beta(s)$ ,  $\lambda(s)$ ,  $k(s)$ ,  $\eta(s)$ ,  $\mu(s)$  are constants the Eqs. (1) and (2) reduce to the corresponding equations in the usual linear thermoelasticity theory.

In the absence of body forces the equation of motion in deformable materials with memory becomes, by the use of Eq. (2),

$$\begin{aligned} \mu(0)\nabla^2\vec{u}(\vec{x}, t) + \{\lambda(0) + \mu(0)\}\text{grad div}\vec{u}(\vec{x}, t) - \beta(0)[\text{grad}\vec{\theta}(\vec{x}, t) + t_1\text{grad}\dot{\vec{\theta}}(\vec{x}, t)] \\ + \int_0^\infty [\mu'(s)\nabla^2\vec{u}(\vec{x}, t-s) + \{\lambda'(s) + \mu'(s)\}\text{grad div}\vec{u}(\vec{x}, t-s) \\ - \beta'(s)[\text{grad}\vec{\theta}(\vec{x}, t-s) + t_1\text{grad}\dot{\vec{\theta}}(\vec{x}, t-s)]]ds = \rho\ddot{\vec{u}}(\vec{x}, t) \end{aligned} \quad (5)$$

Taking the divergence of the Eq. (5), we obtain

$$\begin{aligned} \{\lambda(0) + 2\mu(0)\}\nabla^2 e(\vec{x}, t) - \beta(0)\left[\nabla^2\theta(\vec{x}, t) + t_1\frac{\partial}{\partial t}\nabla^2\theta(\vec{x}, t)\right] \\ + \int_0^\infty \left\{\lambda'(s) + 2\mu'(s)\right\}\nabla^2 e(\vec{x}, t-s) - \beta'(s)\left\{\nabla^2\theta(\vec{x}, t-s) + t_1\frac{\partial}{\partial t}\nabla^2\theta(\vec{x}, t-s)\right\}ds = \rho\ddot{e}(\vec{x}, t) \end{aligned} \quad (6)$$

where  $e(\vec{x}, t) = \text{div}\vec{u}(\vec{x}, t)$ .

### 3. Solution of the problem

Let us assume the following form of the plane wave solutions of Eqs. (1) and (6),

$$[e(\vec{x}, t), \theta(\vec{x}, t)] = (e_0, \theta_0) \exp[i\omega(t - \vec{x} \cdot \hat{n}/c)] \quad (7)$$

where  $\omega > 0$  and  $\hat{n}$  is a unit vector.

Substituting Eq. (7) in Eqs. (6) and (1), we get

$$[\{\lambda(0) + 2\mu(0)\} + \{\bar{\lambda}'(\omega) + 2\bar{\mu}'(\omega)\} - \rho c^2]e_0 - [\beta(0)(1 + t_1 i\omega) + \bar{\beta}'(\omega)\{1 + t_1 i\omega\}]\theta_0 = 0 \quad (8a)$$

$$[\alpha(0)(i\omega - t_0 \omega^2) + \bar{\alpha}'(\omega)\{i\omega - t_0 \omega^2\}]\theta_0 + [(i\omega) + \delta_{lk} t_0 (-\omega^2)]\eta(0)e_0 \\ + e_0[i\omega + t_0 \delta_{lk} (-\omega^2)]\bar{\eta}'(\omega) = [k(0) + \bar{k}'(\omega)](-\omega^2/c^2)\theta_0$$

or,

$$[(i\omega - t_0 \omega^2)\{\gamma(0) + \bar{\gamma}'(\omega)\} + (\omega^2/c^2)\{k(0) + \bar{k}'(\omega)\}]\theta_0 \\ + (i\omega - \delta_{lk} t_0 \omega^2)\{\eta(0) + \bar{\eta}'(\omega)\}e_0 = 0 \quad (8b)$$

where

$$(\bar{\alpha}'(\omega), \bar{\beta}'(\omega), \bar{\gamma}'(\omega), \bar{\lambda}'(\omega), \bar{\mu}'(\omega), \bar{\eta}'(\omega), \bar{k}'(\omega)) = \\ \int_0^\infty (\alpha'(s), \beta'(s), \gamma'(s), \lambda'(s), \mu'(s), \eta'(s), k'(s)) \exp(-i\omega s) ds \quad (9)$$

Putting

$$\alpha(0) + \bar{\alpha}'(\omega) = A, \quad \beta(0) + \bar{\beta}'(\omega) = B, \quad \lambda(0) + 2\mu(0) + \bar{\lambda}'(\omega) + 2\bar{\mu}'(\omega) = L \\ k(0) + \bar{k}'(\omega) = K, \quad \mu(0) + \bar{\mu}'(\omega) = M, \quad \eta(0) + \bar{\eta}'(\omega) = N \quad (10)$$

and eliminating  $e_0, \theta_0$  between Eqs. (8a) and (8b), we get

$$(1 + i\omega t_0 \delta_{lk})\rho c^4 A - c^2[(1 + i\omega \delta_{lk} t_0)AL + \rho i\omega K + (1 + t_1 i\omega)(1 + \delta_{lk} t_0 i\omega)BN] + i\omega LK = 0 \quad (11)$$

If  $t_0 = t_1 = 0$ , we get back the results 6(a) and 6(b) of Chakraborty (1976) in classical Thermoelasticity.

The roots  $C_{1,2}^2$  of Eq. (11) are given by

$$C_{1,2}^2 = [AL(1 + i\omega t_0 \delta_{lk}) + \rho i\omega K + (1 + i\omega t_1)(1 + i\omega t_0 \delta_{lk})BN \pm D]/[2\rho A(1 + i\omega t_0 \delta_{lk})] \quad (12)$$

where

$$D^2 = \{AL(1 + i\omega t_0 \delta_{lk}) + \rho i\omega K + (1 + i\omega t_1)(1 + i\omega t_0 \delta_{lk})BN\}^2 - 4i\omega LK(1 + i\omega t_0 \delta_{lk})\rho A \quad (13)$$

If  $\eta(s) \neq 0$ , the root  $C_1$  corresponds to quasi-elastic wave and the root  $C_2$  corresponds to quasi-thermal wave, because for  $\eta(s) = 0$  and  $t_0 = 0$ ,

$$C_1 = (L/\rho)^{1/2}, \quad C_2 = (i\omega K/A)^{1/2}$$

and these roots correspond, in the uncoupled problem, to a pure elastic and thermal wave respectively.

Let us now analyse the roots of the dispersion Eq. (11) for low and high frequencies.

### (a) Low frequency

Expanding  $\alpha'(\omega)$  for small  $\omega$  in the form

$$\bar{\alpha}'(\omega) = \bar{\alpha}'(0) + \left[ \frac{d}{d\omega} \bar{\alpha}'(\omega) \right]_{\omega=0} \omega + \dots \quad (14)$$

we can write

$$\bar{\alpha}'(\omega) \approx \alpha(\infty) - \alpha(0) - i\alpha_1 \omega$$

where

$$\alpha_1 = \int_0^\infty s \alpha'(s) ds \quad (15)$$

If we assume the existence of integrals of the type

$$\alpha_n = \int_0^\infty s^n \alpha^{(n)}(s) ds, \quad n > 1$$

we can retain terms of higher order in  $\omega$  in Eq. (14).

Thus, for small  $\omega$ , we can write

$$\begin{aligned} A &\approx \alpha(\infty) - i\alpha_1 \omega, \quad B \approx \beta(\infty) - i\omega\beta_1, \quad M \approx (\mu(\infty) - i\mu_1 \omega) \\ L &\approx \lambda(\infty) + 2\mu(\infty) - i\omega(\lambda_1 + 2\mu_1) = l(\infty) - i\omega l_1 \\ K &\approx k(\infty) - i\omega k_1, \quad N \approx \eta(\infty) - i\omega \eta_1 \end{aligned} \quad (16)$$

where  $\alpha_1, \beta_1, \lambda_1, \mu_1, \eta_1, k_1$  are integrals, assumed to converge, of the type Eq. (15). Using the results Eq. (16) in Eq. (12) and extracting the square root, we get for small  $\omega$

$$\begin{aligned} C_1 &\approx \left[ \frac{l(\infty)}{\rho} \right]^{1/2} \cdot \left[ 1 + \frac{\beta(\infty)\eta(\infty)}{l(\infty)\alpha(\infty)} \right]^{1/2} \\ &\times \left[ 1 + i\frac{\omega}{2} \left\{ \frac{t_0 \delta_{lk} l(\infty) \alpha(\infty) - \alpha_1 l(\infty) - l_1 \alpha(\infty) - \beta_1 \eta(\infty) - \eta_1 \beta(\infty) + (t_1 + t_0 \delta_{lk}) \beta(\infty) \eta(\infty)}{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)} \right. \right. \\ &\quad \left. \left. - \frac{t_0 \alpha(\infty) \delta_{lk} - \alpha_1}{\alpha(\infty)} + \rho \frac{k(\infty) \beta(\infty) \eta(\infty)}{(l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty))^2} \right\} \right] \quad (17) \end{aligned}$$

$$\begin{aligned} C_2 &\approx \left[ \frac{\omega k(\infty)}{2l(\infty)} \right]^{1/2} \cdot \left[ 1 + \frac{\beta(\infty)\eta(\infty)}{l(\infty)\alpha(\infty)} \right]^{-1/2} \\ &\times \left[ \left\{ 1 + \frac{\omega}{2l(\infty)} (t_0 \alpha(\infty) \delta_{lk} - \alpha_1) \right\} + i \left\{ 1 - \frac{\omega}{2l(\infty)} (t_0 \alpha(\infty) \delta_{lk} - \alpha_1) \right\} \right] \quad (18) \end{aligned}$$

If  $V_1, V_2$  denote the wave velocities and  $\nu_1, \nu_2$  the attenuation coefficients for the modified elastic and thermal waves respectively, then

$$V_i = \frac{|C_i|^2}{\operatorname{Re} C_i}, \quad i = 1, 2 \quad (19)$$

$$\nu_i = \omega \frac{\operatorname{Im} C_i}{|C_i|^2}, \quad i = 1, 2 \quad (20)$$

Form Eqs. (17)-(20), we obtain

$$V_1 \approx \left[ \frac{\lambda(\infty) + 2\mu(\infty)}{\rho} \right]^{1/2} \left[ 1 + \frac{\beta(\infty)\eta(\infty)}{l(\infty)\alpha(\infty)} \right]^{1/2} [1 + O(\omega^2)] \quad (21)$$

$$\nu_1 \approx -\frac{\omega^2}{2} \left[ \frac{\rho}{\lambda(\infty) + 2\mu(\infty)} \right]^{1/2} \left[ 1 + \frac{\beta(\infty)\eta(\infty)}{l(\infty)\alpha(\infty)} \right]^{-1/2} \quad (22)$$

$$\times \left[ \frac{\alpha_1 l(\infty) + l_1 \alpha(\infty) + \beta_1 \eta(\infty) + \eta_1 \beta(\infty) - t_0 \delta_{lk} l(\infty) \alpha(\infty) - (t_1 + t_0 \delta_{lk}) \beta(\infty) \eta(\infty)}{l(\infty) \alpha(\infty) + \eta(\infty) \beta(\infty)} \right. \\ \left. - \frac{l_1 - \alpha(\infty) \delta_{lk} t_0}{\alpha(\infty)} - \rho \frac{k(\infty) \beta(\infty) \eta(\infty)}{(l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty))^2} \right] \\ V_2 \approx \left[ \frac{2\omega k(\infty)}{l(\infty)} \right]^{1/2} \left[ 1 + \frac{\beta(\infty)\eta(\infty)}{l(\infty)\alpha(\infty)} \right]^{-1/2} \left[ 1 - \frac{\omega}{2} \left\{ \frac{\alpha(\infty) t_0 \delta_{lk} - \alpha_1}{l(\infty)} \right\} \right] \quad (23)$$

$$\nu_2 \approx \left[ \frac{\omega l(\infty)}{2k(\infty)} \right]^{1/2} \left[ 1 + \frac{\beta(\infty)\eta(\infty)}{l(\infty)\alpha(\infty)} \right]^{1/2} \left[ 1 - \frac{\omega}{2} \left\{ \frac{\alpha(\infty) t_0 \delta_{lk} - \alpha_1}{l(\infty)} \right\} \right] \quad (24)$$

From Eq. (21) we see that no relaxation parameter is present, if we neglect terms of higher order of  $\omega$  than the first. But, if we consider terms up to  $\omega^2$ , then  $V_1$  contains terms involving  $t_0$  and  $t_1$ . The above results completely agrees with Chakraborty (1976), if we take  $t_0 = t_1 = 0$

### (b) High frequency

Integrating each of the transforms in Eq. (9) by parts, we obtain

$$\bar{\alpha}'(\omega) = -\frac{\alpha'(0)}{\omega} i - \frac{\alpha''(0)}{\omega^2} - \frac{\bar{\alpha}'''(\omega)}{\omega^2} \quad (25)$$

$$\bar{\beta}'(\omega) = -\frac{\beta'(0)}{\omega} i - \frac{\beta''(0)}{\omega^2} - \frac{\bar{\beta}'''(\omega)}{\omega^2} \quad \text{etc.} \quad (26)$$

where

$$\bar{\alpha}'''(\omega) = \int_0^\infty \alpha'''(s) \exp(-i\omega s) ds, \quad \text{etc.}$$

Hence for large  $\omega$ , we can write

$$\begin{aligned} A &\approx \alpha(0) - \frac{i}{\omega} \alpha'(0), \quad B \approx \beta(0) - \frac{i}{\omega} \beta'(0) \\ K &\approx k(0) - \frac{i}{\omega} k'(0), \quad L \approx (\lambda(0) + 2\mu(0)) - \frac{i}{\omega} (\lambda'(0) + 2\mu'(0)) \\ M &\approx \mu(0) - \frac{i}{\omega} \mu'(0), \quad N \approx \eta(0) - \frac{i}{\omega} \eta'(0) \end{aligned} \quad (27)$$

The substitution of Eq. (27) in Eq. (12) yields, for large  $\omega$ ,

$$\begin{aligned} C_1 &\approx \left[ \frac{\omega k(0)}{2\alpha(0)} \right]^{1/2} \left[ 1 + \frac{1}{2\omega} \left\{ \frac{\beta(0)\eta(0)}{\rho k(0)} + \frac{k'(0)}{k(0)} - \frac{\alpha'(0)}{\alpha(0)} + \frac{1}{\rho k(0)} (t_1 + t_0 \delta_{lk}) \beta(0)\eta(0) \right\} \right. \\ &\quad \left. + i \left\{ 1 - \frac{1}{2\omega} \left( \frac{\beta(0)\eta(0)}{\rho k(0)} + \frac{k'(0)}{k(0)} - \frac{\alpha'(0)}{\alpha(0)} + \frac{1}{\rho k(0)} (t_1 + t_0 \delta_{lk}) \beta(0)\eta(0) \right) \right\} \right] \end{aligned} \quad (28)$$

$$C_2 \approx \left[ \frac{\lambda(0) + 2\mu(0)}{\rho} \right]^{1/2} \left[ 1 - \left( \frac{i}{2\omega} \cdot \frac{\lambda'(0) + 2\mu'(0)}{\lambda(0) + 2\mu(0)} \right) \right] \quad (29)$$

The wave velocities and attenuation coefficients for the elastic and thermal waves are given by

$$V_1 \approx \left[ \frac{2\omega k(0)}{\alpha(0)} \right]^{1/2} \left[ 1 - \frac{1}{2\omega} \left\{ \frac{\beta(0)\eta(0)}{\rho k(0)} + \frac{k'(0)}{k(0)} - \frac{\alpha'(0)}{\alpha(0)} + \frac{1}{\rho k(0)} (t_1 + t_0 \delta_{lk}) \beta(0)\eta(0) \right\} \right] \quad (30)$$

$$\nu_1 \approx \left[ \frac{\omega \alpha(0)}{2k(0)} \right]^{1/2} \left[ 1 - \frac{1}{2\omega} \left\{ \frac{\beta(0)\eta(0)}{\rho k(0)} + \frac{k'(0)}{k(0)} - \frac{\alpha'(0)}{\alpha(0)} + \frac{1}{\rho k(0)} (t_1 + t_0 \delta_{lk}) \beta(0)\eta(0) \right\} \right] \quad (31)$$

$$V_2 \approx \left[ \frac{\lambda(0) + 2\mu(0)}{\rho} \right]^{1/2} \left[ 1 + 0 \left( \frac{1}{\omega^2} \right) \right] \quad (32)$$

$$\nu_2 \approx -\frac{1}{2} \left( \frac{\rho}{\lambda(0) + 2\mu(0)} \right)^{1/2} \left[ \frac{\lambda'(0) + 2\mu'(0)}{\lambda(0) + 2\mu(0)} \right] \left[ 1 + 0 \left( \frac{1}{\omega^2} \right) \right] \quad (33)$$

It is observed that both the modified elastic and thermal waves in generalized thermoelastic medium exhibit damping and dispersion. The modified thermal elastic wave speed, unlike that in the classical result approaches the instantaneous value of the classical compressional wave velocity at high frequencies and its equilibrium value is modified due to the presence of the coupling factor at low frequencies. Both the quasi-thermal wave speed and attenuation coefficient are proportional to the square root of the frequency at low frequency. Results for 'thermoelastic' materials may be obtained from the results Eqs. (21)-(24) and Eqs. (30)-(33) by taking the relaxation functions to constants. They agree with those given by Chadwik (1960) and Nowacki (1962).

We neglect the coupling between the temperature and the displacement fields by putting



$$\eta(s) = \eta(0) = \eta(\infty) = 0$$

in Eqs. (21)-(24) and Eqs. (30)-(33) and get the corresponding results for the purely elastic and the purely thermal modes. Results for purely thermal mode agree with those given by Nunziato (1971).

For the purely elastic mode

$$V_E \approx \left[ \frac{\lambda(\infty) + 2\mu(\infty)}{\rho} \right]^{1/2} \left[ 1 + \frac{\omega^2}{4} \left( \frac{\lambda_1 + 2\mu_1}{\lambda(\infty) + 2\mu(\infty)} \right)^2 \right] \quad \text{as } \omega \rightarrow 0 \quad (34)$$

$$\nu_E \approx -\frac{\omega^2}{2} \left[ \frac{\rho}{\lambda(\infty) + 2\mu(\infty)} \right]^{1/2} \left[ \frac{\lambda_1 + 2\mu_1}{\lambda(\infty) + 2\mu(\infty)} \right] \quad \text{as } \omega \rightarrow 0 \quad (35)$$

where  $V_E$  and  $\nu_E$  are the wave speed and attenuation coefficient for the purely elastic waves. Also,

$$V_E \approx \left[ \frac{\lambda(0) + 2\mu(0)}{\rho} \right]^{1/2} \left[ 1 + \frac{1}{4} \omega^2 \left( \frac{\lambda_1 + 2\mu_1}{\lambda(0) + 2\mu(0)} \right)^2 \right] \quad \text{as } \omega \rightarrow \infty \quad (36)$$

$$\nu_E \approx -\frac{1}{2} \left( \frac{\rho}{\lambda(0) + 2\mu(0)} \right)^{1/2} \left[ \frac{\lambda'(0) + 2\mu'(0)}{\lambda(0) + 2\mu(0)} \right] \left[ 1 + 0 \left( \frac{1}{\omega^2} \right) \right] \quad \text{as } \omega \rightarrow \infty \quad (37)$$

The results Eqs. (34)-(37) show that the purely elastic wave has its modification, and the quasi-elastic wave is subject to damping and dispersion. This is a feature exhibited by the elastic waves only in material with memory both in the cases of classical and generalized thermoelasticity. The attenuation coefficients at low and high frequencies are proportional to  $\lambda_1 + 2\mu_1$  and  $\lambda'(0) + 2\mu'(0)$  respectively. These will vanish if  $\lambda(s) + 2\mu(s)$  is a constant i.e., in classical elasticity. Thus attenuation of pure elastic wave occurs in materials with memory both in the cases of classical and generalized thermoelasticity. However, it may be pointed out that the effect of generalized thermoelasticity is found to be absent. The velocity for pure elastic waves tends to different limits, in general, according as  $\omega \rightarrow 0$  or  $\omega \rightarrow \infty$ ; but if  $\lambda(s) + 2\mu(s)$  is a constant, as in classical elasticity  $V_E$  becomes independent of  $\omega$  and assumes the constant value of the classical compressional wave velocity.

#### 4. Rayleigh waves

Putting the displacement vector  $\vec{u}$  in the form

$$\vec{u} = \text{grad}\phi + \text{rot}\vec{A}, \quad \vec{A} = (0, 0, \psi)$$

as the sum of irrotational and solenoidal components in the Eqs. (1) and (5), we get

$$\begin{aligned} & \left[ \{ \lambda(0) + 2\mu(0) \} \nabla^2 - \rho \frac{\partial^2}{\partial t^2} \right] \phi(\vec{x}, t) - \beta(0) [\theta(\vec{x}, t) + t_1 \dot{\theta}(\vec{x}, t)] \\ & + \int_0^\infty [\{ \lambda'(s) + 2\mu'(s) \} \nabla^2 \phi(\vec{x}, t-s) - \beta'(s) \{ \theta(\vec{x}, t) + t_1 \dot{\theta}(\vec{x}, t) \}] ds = 0 \end{aligned} \quad (38)$$

$$\mu(0)\nabla^2\psi(\vec{x},t) - \rho\ddot{\psi}(\vec{x},t) + \int_0^\infty \mu'(s)\nabla^2\psi(\vec{x},t-s)ds = 0 \quad (39)$$

$$\begin{aligned} & \alpha(0)(\dot{\theta} + t_0\ddot{\theta}) + \int_0^\infty \gamma'(s)[\dot{\theta}(\vec{x},t-s) + t_0\ddot{\theta}(\vec{x},t-s)]ds + \eta(0)\{\nabla^2\dot{\phi}(\vec{x},t) + \delta_{lk}t_0\nabla^2\ddot{\phi}(\vec{x},t)\} \\ & + \int_0^\infty \eta'(s)\{\nabla^2\dot{\phi}(\vec{x},t-s) + \delta_{lk}t_0\ddot{\phi}(\vec{x},t-s)\}ds = k(0)\nabla^2\theta(\vec{x},t) + \int_0^\infty k'(s)\nabla^2\theta(\vec{x},t-s)ds \end{aligned} \quad (40)$$

Using the substitution (Nowacki 1962)

$$\{\theta(\vec{x},t), \phi(\vec{x},t), \psi(\vec{x},t)\} = \{\hat{\theta}(x_1), \hat{\phi}(x_1), \hat{\psi}(x_1)\}\exp[i\omega(t-x_2/C)]$$

and Eq. (10) in Eqs. (38)-(40) we get

$$\begin{aligned} & [KL(D^2 - \omega^2/C^2)^2 - i\omega(D^2 - \omega^2/C^2)\{(1+it_0\omega)AL + (1+i\omega t_1)(1+it_0\omega)BN + i\rho\omega K\} \\ & - i\omega^3\rho(1+it_0\omega)A](\hat{\theta}, \hat{\phi}) = 0 \end{aligned} \quad (41)$$

$$[M(D^2 - \omega^2/C^2) + \rho\omega^2]\hat{\psi} = 0 \quad (42)$$

where

$$D \equiv \frac{d}{dx_1}$$

For surface waves we assume the solutions of Eqs. (41)-(42) in the form

$$\hat{\psi}(x_1) = A_3\exp(-\gamma x_1) \quad (43)$$

and

$$\hat{\phi}(x_1) = A_1\exp(-x_1\chi_1) + A_2\exp(-x_1\chi_2) \quad (44)$$

where  $\chi_1$  and  $\chi_2$  are the roots of the equation

$$\begin{aligned} & [KL(\chi^2 - a^2)^2 - i\omega(\chi^2 - a^2)\{(1+it_0\omega)AL + (1+i\omega t_1)(1+i\omega t_0)BN + i\rho\omega K\} \\ & - i\omega^3\rho(1+t_0i\omega)A] = 0 \end{aligned} \quad (45)$$

If  $t_0$  and  $t_1$  are each equal to zero, the Eq. (45) totally coincides with that obtained by Chakraborti (1976). Now putting the expression Eq. (44) in Eq. (38), we get

$$\begin{aligned} \hat{\theta}_1(x_1) = & \frac{1}{B(1+i\omega t_1)}[\{(L(\chi_1^2 - a^2) + \rho\omega^2)A_1\exp(-\chi_1 x_1) \\ & + (L(\chi_2^2 - a^2) + \rho\omega^2)A_2\exp(-\chi_2 x_1)\}] \end{aligned} \quad (46)$$

The stress components  $\sigma_{11}(\vec{x},t)$  and  $\sigma_{12}(\vec{x},t)$  given by Eq. (2) can be rewritten in terms of the

potential functions  $\phi$ ,  $\psi$  as

$$\begin{aligned}\sigma_{12} = & 2\mu(0)\phi_{,12}(\vec{x}, t) + 2 \int_0^\infty \mu'(s)\phi_{,12}(\vec{x}, t-s)ds \\ & + \mu(0)[\psi_{,22}(\vec{x}, t) - \psi_{,11}(\vec{x}, t)] + \int_0^\infty \mu'(s)[\psi_{,22}(\vec{x}, t-s) - \psi_{,11}(\vec{x}, t-s)]ds\end{aligned}\quad (47)$$

$$\begin{aligned}\sigma_{11}(\vec{x}, t) = & \rho\ddot{\phi}(\vec{x}, t) + 2\mu(0)[\psi_{,12}(\vec{x}, t) - \phi_{,22}(\vec{x}, t)] \\ & + 2 \int_0^\infty \mu'(s)[\psi_{,12}(\vec{x}, t-s) - \phi_{,22}(\vec{x}, t-s)]ds\end{aligned}\quad (48)$$

Then the boundary conditions

$$\begin{aligned}\frac{\partial\theta(0, x_2; t)}{\partial x_1} + h\theta(0, x_2; t) &= 0 \\ \sigma_{11}(0, x_2; t) &= 0 = \sigma_{12}(0, x_2; t)\end{aligned}$$

will yield the system of equations

$$\begin{aligned}(2a^2 - \tau^2)A_1 + (2a^2 - \tau^2)A_2 + 2ia\gamma A_3 &= 0 \\ 2ia\chi_1 A_1 + 2ia\chi_2 A_2 - (2a^2 - \tau^2)A_3 &= 0 \\ (h - \chi_1)n_1 A_1 + (h - \chi_2)n_2 A_2 &= 0\end{aligned}\quad (49)$$

where  $n_{1,2} = \chi_{1,2}^2 + \sigma^2 - a^2$ ,  $\sigma^2 = \rho\omega^2/L$ ,  $a^2 = \omega^2/C^2$ ,  $\tau = \rho\omega^2/M$ ,  $\gamma = (a^2 - \tau^2)^{1/2}$

The condition of consistency of the system of Eq. (49) gives

$$(2a^2 - \tau^2)^2 = 4a^2(a^2 - \tau^2)^{1/2} \frac{(\chi_1 n_2 - \chi_2 n_1)h + \chi_1 \chi_2 (n_1 - n_2)}{(n_1 - n_2)h + (\chi_1 n_1 - \chi_2 n_2)}$$

$$\begin{aligned}\text{i.e., } (2a^2 - \tau^2)^2 [(\sigma^2 - a^2) + \chi_1^2 + \chi_2^2 + \chi_1 \chi_2] - 4a^2(a^2 - \tau^2)^{1/2} \chi_1 \chi_2 (\chi_1 + \chi_2) \\ = h[(2a^2 - \tau^2)^2 (\chi_1 + \chi_2) + 4a^2(a^2 - \tau^2)^{1/2} \{(\sigma^2 - a^2) - \chi_1 \chi_2\}]\end{aligned}\quad (50)$$

For small  $\omega$ ,  $\omega \ll 1$ , this Eq. (50) becomes [see Appendix]

$$\begin{aligned}\left[ i + i^{1/2} \omega^{1/2} \left\{ \frac{1}{C^2} - \frac{\rho\alpha(\infty)}{l(\infty)\alpha(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \cdot \left\{ \frac{l(\infty)\alpha(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{-1/2} \right] \\ \times \left[ \left( \frac{2}{C^2} - \frac{\rho}{\mu(\infty)} \right)^2 - \frac{4}{C^2} \left( \frac{1}{C^2} - \frac{\rho}{\mu(\infty)} \right)^{1/2} \cdot \left\{ \frac{1}{C^2} - \frac{\rho\mu(\infty)}{l(\infty)\alpha(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right]\end{aligned}$$

$$\begin{aligned}
&= \frac{h}{\omega^{1/2}} \left\{ \left( \frac{2}{C^2} - \frac{\rho}{\mu(\infty)} \right)^2 \cdot \left[ i^{1/2} + \omega^{1/2} \left\{ \frac{l(\infty)\alpha(\infty) + \beta(\infty)\eta(\infty)}{\alpha(\infty)k(\infty)} \right\} \right]^{-1/2} \right. \\
&\quad \times \left. \left\{ \frac{1}{C^2} - \frac{\rho\alpha(\infty)}{l(\infty)\alpha(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right] - \frac{4}{C^2} \left( \frac{1}{C^2} - \frac{\rho}{\mu(\infty)} \right)^{1/2} \cdot i^{1/2} \\
&\quad \left[ \frac{1}{C^2} - \frac{\rho\alpha(\infty)}{l(\infty)\alpha(\infty) + \beta(\infty)\eta(\infty)} \right]^{1/2} \left\{ \frac{l(\infty)\alpha(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{-1/2} [1 + o(\omega^{1/2})] \quad (51)
\end{aligned}$$

The result Eq. (51) shows that, on neglecting terms of order  $\omega^{1/2}$ , the velocity  $C$  ceases to depend upon the frequency  $\omega$  and the thermal constant  $h$  (Chadwick 1960).

We then have

$$\left[ 2 - \frac{\rho C^2}{\mu(\infty)} \right]^2 = 4 \left[ 1 - \frac{\rho C^2}{\mu(\infty)} \right]^{1/2} \cdot \left[ 1 - \frac{\rho C^2 \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right]^{1/2}$$

This result is analogous to the corresponding result in classical thermoelasticity.

## 5. Conclusions

It is observed that both the modified elastic and thermal waves in generalized thermoelastic medium exhibit damping and dispersion. The modified thermal elastic wave speed, unlike that in the classical result approaches the instantaneous value of the classical compressional wave velocity at high frequencies and its equilibrium value is modified due to the presence of the coupling factor at low frequencies. Both the quasi-thermal wave speed and attenuation coefficient are proportional to the square root of the frequency at low frequency.

## Acknowledgements

We express our sincere thanks to the learned reviewer for his comments and suggestions for the improvement of the paper.

## References

- Amos, D.E. and Chen, P.J. (1970), *J. Appl. Mech.* **37**, Series E, No. 4, 1145-1146.
- Bhattacharyya, R.K. (1991), Wave Propagation in Random Thermoelastic Materials with Memory, in Mathematical and Numerical Aspects of Wave Propagation Phenomena, (Strasbourg 1991), 626-634, SIMA, Philadelphia, PA, 1991; secMR92a:65021; CNO 1106033.
- Chadwick, P. (1960), Thermoelasticity. The Dynamical Theory. Progress in Solid Mech. (Edited by Sneddon and Hill) Vol. 1, North- Holland Publishing Co. Amsterdam, 307-311.
- Chakraborti, S. (1976), "Waves in thermoelastic materials with memory", *Bull. Math. De la R. S. de Roumanie Tome*, **20**(68), nr. 1-2.
- Chen, P.J. (1969), "On the growth and decay of temperature rate waves of arbitrary form", *ZAMP*, **20**, 448-453.

- Chen, P.J. and Gurtin, M.E. (1970), "On second sound in materials with memory", *ZAMP*, **21**, 232-241.
- Chirita, S. and Quintanilla, R. (1997), "Spatial estimates in the dynamic theory of linear elastic materials with memory", *European J. Mech./A Solids*, **16**, 723-736.
- Dhaliwal, R.S. and Singh, A. (1980), *Dynamic Coupled Thermoelasticity*, Hindustan Publ., Delhi.
- Eringen, A.C. (1967), *Mechanics of Continua*, John Wiley and Sons, IUNC, New York.
- Gao, Hongjun. (2001), "On the exponential stability of thermoelastic problem with memory", *Applicable Anal. an Int. J.*, **78**, Mo. 3-4, 379-403.
- Green, A.E. and Lindsay, K.A. (1972), Thermoelasticity, *J. Elasticity*, **2**, 1-7.
- Gurtin, M.E. and Pipkin, A.C. (1968), "A general theory of heat conduction with finite wave speed", *Arch. Rat. Mech. Anal.*, **31**, 113-126.
- Iesan, D. and Quintanilla, R. (2002), "On a theory of interacting continua with memory", *J. Thermal Stresses*, **25**, 1161-1178.
- Lord, H.W. and Shulman, Y. (1967), "A generalized dynamical theory of thermoelasticity", *J. Mech. Phys. Solids*, **15**, 299-309.
- Matinez, F. and Quintanilla, R. (1995), "Existenciay unicidad de soluciones para las ecuaciones de la termoelasticidad de materials con memoria", Actas 111 Congreso de Matematica Aplicada. (Ed. Casal *et al.*).
- McCarthy, M. (1970), *Proc. Vibration Problems*, **11**(2), 123- 133.
- McCarthy, M. (1970), "Constitutive equations for thermomechanical materials with memory", *Int. J. Eng. Sci.*, **8**, 467.
- Müller, I. (1967), *Arch. Rat. Mech. Anal*, 118-141.
- Naso. Maria Garzia (2003), "Exact boundary controllability in thermoelasticity with memory", *Advan. Differential Equations*, **8**(4), 471-490.
- Naso. Maria Garzia and Vegni. Federico. (2003), "Asymptotic behavior for second-order systems with memory", *J. Math. Anal. Appl.*, **286**(2), 692-794.
- Naso. Maria Garzia and Vuk, Elena. (2004), "Asymptotic behaviour of the energy for electromagnetic system with memory", *J. Math. Meth. Appl. Sci.*, **27**(7), 819-841.
- Naso. Maria Garzia and Yuming. Qin. (2002), "Global existence exponential stability in one-dimensional nonlinear thermoelasticity with thermal memory", *Non Linear Anal.*, **51**, No. 1-A, 11-32.
- Nowacki, W. (1962), *Thermoelasticity*, Pergamon Press, 298.
- Nunziato, J.W. (1971), "On heat conduction in materials with memory", *Q. Appl. Mech.*, **29**, 187-204.
- Zhou, W.M., Liu, Y., Jiang, B.H., Qi, X. and Liu, Y.N. (2003), "Magnetoelastic and thermoelastic shape memory effect in a Co-Ni single crystal", *Applied Phys. Letters*, **82**(5), 760-762.

## Appendix

$\chi_1^2$  and  $\chi_2^2$  are the roots of the equation

$$[KL(\chi^2 - a^2)^2 - i\omega(\chi^2 - a^2)\{(1 + it_0\omega)AL + (1 + it_1\omega)(1 + it_0\omega)BN + i\rho\omega K\} - i\omega^3\rho(1 + t_0i\omega)A] = 0$$

or

$$LK\chi^4 - \chi^2[2a^2LK + i\omega\{(1 + it_0\omega)AL + (1 + it_1\omega)(1 + it_0\omega)BN + i\rho\omega K\}] + [a^4LK + i\omega a^2\{(1 + it_0\omega)AL + (1 + it_1\omega)(1 + it_0\omega)BN + i\rho\omega K\} - i\omega^3\rho(1 + it_0\omega)A] = 0$$

Therefore

$$\chi_1^2 + \chi_2^2 = [2a^2LK + i\omega\{(1 + it_0\omega)AL + (1 + it_1\omega)(1 + it_0\omega)BN + i\rho\omega K\}]/LK$$

and

$$\chi_1^2 \cdot \chi_2^2 = [a^4LK + i\omega a^2\{(1 + it_0\omega)AL + (1 + it_1\omega)(1 + it_0\omega)BN + i\rho\omega K\} - i\omega^3\rho(1 + it_0\omega)A]/LK$$

Now  $a^2 = \omega^2/C^2$

$$A \approx \alpha(\infty) - i\alpha_1\omega, \quad B \approx \beta(\infty) - i\omega\beta_1, \quad M \approx (\mu(\infty) - i\mu_1\omega)$$

$$L \approx \lambda(\infty) + 2\mu(\infty) - i\omega(\lambda_1 + 2\mu_1) = l(\infty) - i\omega l_1$$

$$K \approx k(\infty) - i\omega k_1, \quad N \approx \eta(\infty) - i\omega \eta_1.$$

$$\chi_1^2 + \chi_2^2 \approx i\omega \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}.$$

$$\chi_1^2 \cdot \chi_2^2 \approx \frac{i\omega^3}{C^2} \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\} \cdot \left\{ 1 - \frac{\rho C^2 \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}.$$

$$\chi_1 \cdot \chi_2 \approx \frac{\sqrt{i}\omega^{3/2}}{C} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{1/2} \cdot \left\{ 1 - \frac{\rho C^2 \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2}.$$

$$(\chi_1 + \chi_2)^2 \approx i\omega \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\} +$$

$$2\frac{\sqrt{i}\omega^{3/2}}{C} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{1/2} \cdot \left\{ 1 - \frac{\rho C^2 \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2}$$

$$= i\omega \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}.$$

$$\left[ 1 + 2\frac{\omega^{1/2}}{i^{1/2}} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{-1/2} \cdot \left\{ \frac{1}{C^2} - \frac{\rho\alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right].$$

$$(\chi_1 + \chi_2) \approx (i\omega)^{1/2} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{1/2}.$$

$$\left[ 1 + \frac{\omega^{1/2}}{i^{1/2}} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{-1/2} \cdot \left\{ \frac{1}{C^2} - \frac{\rho\alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right].$$

$$(2a^2 - \tau^2) \approx \omega^4 \left( \frac{2}{C^2} - \frac{\rho}{\mu(\infty)} \right)^2.$$

$$(\sigma^2 - a^2) + \chi_1^2 + \chi_2^2 + \chi_1\chi_2 \approx i\omega \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\} \times$$

$$\left[ 1 + \frac{\omega^{1/2}}{i^{1/2}} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{-1/2} \cdot \left\{ \frac{1}{C^2} - \frac{\rho\alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right]$$

$$4a^2(\sigma^2 - a^2)^{1/2} \chi_1\chi_2(\chi_1 + \chi_2) \approx$$

$$\begin{aligned}
& \frac{4i\omega^5}{C^2} \left( \frac{1}{C^2} - \frac{\rho}{\mu(\infty)} \right)^{1/2} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{1/2} \cdot \left\{ 1 - \frac{\rho c^2 \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \times \\
& \left[ 1 + \frac{\omega^{1/2}}{i^{1/2}} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{-1/2} \cdot \left\{ \frac{1}{C^2} - \frac{\rho \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right] \cdot \\
& (2a^2 - \tau^2)(\chi_1 + \chi_2) \approx \omega^4 (\omega i)^{1/2} \left( \frac{1}{C^2} - \frac{\rho}{\mu(\infty)} \right)^2 \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{1/2} \times \\
& \left[ 1 + \frac{\omega^{1/2}}{i^{1/2}} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{-1/2} \cdot \left\{ \frac{1}{C^2} - \frac{\rho \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right] \cdot \\
& 4a^2 (\sigma^2 - a^2)^{1/2} \{ (\sigma^2 - a^2) - \chi_1 \chi_2 \} \approx \frac{4\omega^5}{C^2} \left( \frac{1}{C^2} - \frac{\rho}{\mu(\infty)} \right)^{1/2} \times \\
& \left[ \left( \frac{\rho}{l(\infty)} - \frac{1}{C^2} \right) - \frac{i^{1/2}}{\omega^{1/2}} \cdot \left\{ \frac{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)}{l(\infty)k(\infty)} \right\}^{1/2} \cdot \left\{ \frac{1}{C^2} - \frac{\rho \alpha(\infty)}{\alpha(\infty)l(\infty) + \beta(\infty)\eta(\infty)} \right\}^{1/2} \right].
\end{aligned}$$

Therefore

$$\begin{aligned}
& (2a^2 - \tau^2)^2 [(\sigma^2 - a^2) + \chi_1^2 + \chi_2^2 + \chi_1 \chi_2] - 4a^2 (\sigma^2 - \tau^2)^{1/2} \chi_1 \chi_2 (\chi_1 + \chi_2) = \\
& h[(2a^2 - \tau^2)^2 (\chi_1 + \chi_2) + 4a^2 (\sigma^2 - \tau^2)^{1/2} \{ (\sigma^2 - a^2) - \chi_1 \chi_2 \}]
\end{aligned}$$

becomes

$$\begin{aligned}
& \left[ 1 + i^{1/2} \omega^{1/2} \left\{ \frac{1}{C^2} - \frac{\rho \alpha(\infty)}{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)} \right\}^{1/2} \cdot \left\{ \frac{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)}{l(\infty) k(\infty)} \right\}^{-1/2} \right] \times \\
& \left[ \left( \frac{2}{C^2} - \frac{\rho}{\mu(\infty)} \right)^2 - \frac{4}{C^2} \left( \frac{1}{C^2} - \frac{\rho}{\mu(\infty)} \right)^{1/2} \frac{1}{C^2} - \frac{\rho \mu(\infty)}{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)} \right]^{1/2} = \frac{h}{\omega^{1/2}} \left\{ \left( \frac{2}{C^2} - \frac{\rho}{\mu(\infty)} \right)^2 \right. \\
& \left. \left[ i^{1/2} + \omega^{1/2} \left\{ \frac{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)}{\alpha(\infty) k(\infty)} \right\}^{-1/2} \times \left\{ \frac{1}{C^2} - \frac{\rho \alpha(\infty)}{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)} \right\}^{1/2} \right] - \right. \\
& \left. \frac{4}{C^2} \left( \frac{1}{C^2} - \frac{\rho}{\mu(\infty)} \right)^{1/2} \cdot i^{1/2} \left[ \frac{1}{C^2} - \frac{\rho \alpha(\infty)}{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)} \right]^{1/2} \right\} \cdot \left\{ \frac{l(\infty) \alpha(\infty) + \beta(\infty) \eta(\infty)}{l(\infty) k(\infty)} \right\}^{-1/2} \\
& [1 + O(\omega^{1/2})]
\end{aligned}$$